

Quantum field theoretic model of metastable resonant spin–singlet state of the np pair

A. N. Ivanov ^{*†}, M. Cargnelli [‡], M. Faber [§], H. Fuhrmann [¶],
V. A. Ivanova ^{||}, J. Marton ^{**}, N. I. Troitskaya ^{††}, J. Zmeskal ^{‡‡}

February 5, 2008

*Atominstitut der Österreichischen Universitäten, Arbeitsbereich Kernphysik und
Nukleare Astrophysik, Technische Universität Wien,
Wiedner Hauptstr. 8-10, A-1040 Wien, Österreich
und
Institut für Mittelenergiephysik Österreichische Akademie der Wissenschaften,
Boltzmannngasse 3, A-1090, Wien, Österreich*

Abstract

The np pair in the spin–singlet state is treated as a Cooper np –pair within the extension of the Nambu–Jona–Lasinio model of light nuclei, describing the deuteron as a Cooper np –pair in the spin–triplet state. For the Cooper np –pair in the spin–singlet state we compute the binding energy and express the S–wave scattering length of np scattering in the spin–singlet state in terms of the binding energy. The theoretical value of the S–wave scattering length of np scattering agrees well with the experimental data.

PACS: 11.10.Ef, 21.10.Dr, 21.30.Fe, 25.70.Ef

^{*}E–mail: ivanov@kph.tuwien.ac.at, Tel.: +43–1–58801–14261, Fax: +43–1–58801–14299

[†]Permanent Address: State Polytechnic University, Department of Nuclear Physics, 195251 St. Petersburg, Russian Federation

[‡]E–mail: michael.cargnelli@oeaw.ac.at

[§]E–mail: faber@kph.tuwien.ac.at, Tel.: +43–1–58801–14261, Fax: +43–1–58801–14299

[¶]E–mail: hermann.fuhrmann@oeaw.ac.at

^{||}E–mail: viola@kph.tuwien.ac.at, State Polytechnic University, Department of Nuclear Physics, 195251 St. Petersburg, Russian Federation

^{**}E–mail: johann.marton@oeaw.ac.at

^{††}State Polytechnic University, Department of Nuclear Physics, 195251 St. Petersburg, Russian Federation

^{‡‡}E–mail: johann.zmeskal@oeaw.ac.at

At very low energies the S-wave amplitude of np scattering in the 1S_0 state (the spin-singlet state) can be written as [1, 2]

$$f_{np}(k) = \frac{1}{-\kappa_{np} - i k}, \quad (1)$$

where $1/\kappa_{np} = a_{np}$ can be identified with the S-wave scattering length of np scattering in the spin-singlet state¹ and k is a relative momentum of the np pair. The experimental value of the S-wave scattering lengths of np scattering in the spin-singlet state is equal to $a_{np} = (-23.75 \pm 0.01)$ fm [2].

According to Landau and Lifshitz [1], the S-wave scattering length $a_{np} = (-23.75 \pm 0.01)$ fm of np scattering in the spin-singlet state testifies the existence of the *virtual level* with the binding energy $\varepsilon_{^1S_0} = \kappa_{np}^2/m_N = 1/a_{np}^2 m_N = 0.074$ MeV, where $m_N = 940$ MeV is the nucleon mass.

It is well-known that strong low-energy interactions of the np pair in the 3S_1 state (the spin-triplet state) lead to the existence of the bound state, the deuteron [2] with the binding energy $\varepsilon_D = 2.225$ MeV [2]. The quantum field theoretic approach to the description of the deuteron as the Cooper np pair has been developed within the Nambu–Jona-Lasinio model of light nuclei in Refs. [3]–[7]².

As has been shown in [3]–[7] the NNJL model describes well the low-energy parameters of the deuteron, such as the binding energy, the dipole magnetic and electric quadrupole moments [3], the $\Delta\Delta$ component of the wave function of the deuteron in the ground state [4], the low-energy nuclear reactions with the deuteron of astrophysical interest [5], the asymptotic ratio D/S [6] of the wave function of the deuteron³ and the energy level displacement of the ground state of pionic deuterium [7].

The existence of the metastable resonance of the np pair in the spin-singlet state with a mass in the vicinity of the threshold mass of the np pair and the possibility to treat this metastable state as a Cooper np -pair has been suggested by Ericson [9] (see also [10]).

In this letter we follow a suggestion by Ericson [9] and treat the np pair in the spin-singlet state as a Cooper pair in a way analogous to the NNJL model [3]–[7]. The local interpolating field operator of the np pair in the spin-singlet state we define as follows $S(x) = n^c(x)i\gamma^5 p(x)$ [3], where $n(x)$ and $p(x)$ are the interpolating field operators of the neutron and the proton, respectively. The effective low-energy Lagrangian, unstable with respect to the production of the metastable resonant np pair in the spin-singlet state and containing only the fields of the neutron and the proton, we define as follows [3]

$$\begin{aligned} \mathcal{L}^{np}(x) &= \bar{n}(x)(i\gamma^\mu\partial_\mu - m_N)n(x) + \bar{p}(x)(i\gamma^\mu\partial_\mu - m_N)p(x) \\ &+ \frac{g_S^2}{4m_N^2} [\bar{p}(x)i\gamma^5 n^c(x)][n^c(x)i\gamma^5 p(x)] + \dots, \end{aligned} \quad (2)$$

where ellipses stand for low-energy interactions of the neutron and the proton with other

¹For simplicity we neglect the contribution of the effective range [1, 2].

²We use the abbreviation the NNJL model that means the Nuclear Nambu–Jona-Lasinio model [3].

³The asymptotic ratio D/S of the D-wave component to the S-wave component of the wave function of the deuteron in the ground state has been computed in the NNJL model in agreement with the results obtained by Ericson within the potential model approach [8] and the experimental value, which has been used by Kamionkowski and Bahcall [8] for the calculation of the astrophysical factor for the solar proton burning $p + p \rightarrow d + e^+ + \nu_e$.

fields [3]. The coupling constant g_S is related to the coupling constant g_V of the NNJL model [3]⁴ as $g_S = g_V/\sqrt{3}$ [5]⁵.

In order to introduce the interpolating local field $S(x)$ we should linearize the Lagrangian (2). Following the procedure described in [3] we get

$$\begin{aligned}\mathcal{L}^{\text{np}}(x) &= \bar{n}(x)(i\gamma^\mu\partial_\mu - m_N)n(x) + \bar{p}(x)(i\gamma^\mu\partial_\mu - m_N)p(x) \\ &- 4m_N^2 S^\dagger(x)S(x) + g_S[n^c(x)i\gamma^5 p(x)]S^\dagger(x) + g_S[\bar{p}(x)i\gamma^5 n^c(x)]S(x) + \dots\end{aligned}\quad (3)$$

For the derivation of the effective Lagrangian of the physical S -field we should integrate over nucleon fields [3].

Following the procedure of the integration over the nucleon fields, expounded in [3], we arrive at the effective Lagrangian of the collective field $S(x) = n^c(x)i\gamma^5 p(x)$:

$$\mathcal{L}(x) = \left(1 + \frac{g_S^2}{8\pi^2} J_2(m_N)\right) \partial_\mu S^\dagger(x) \partial^\mu S(x) - \left(4m_N^2 - \frac{g_S^2}{4\pi^2} J_1(m_N)\right) S^\dagger(x) S(x), \quad (4)$$

where $J_1(m_N)$ and $J_2(m_N)$ are divergent integrals related by [3]

$$J_1(m_N) = 2m_N^2 J_2(m_N) = \frac{4}{3} \frac{\Lambda_D^3}{m_N} \sim O(1/N_C). \quad (5)$$

The cut-off $\Lambda_D = 46$ MeV [3] restricts from above 3-momenta of low-energy fluctuations of virtual neutron and proton fields forming the physical metastable resonant np state [3]. Making a renormalization of the wave function of the collective field $S(x)$ [3], $S(x) \rightarrow Z^{1/2} S(x)$, where $Z = 1 + (g_S^2/8\pi^2) J_2(m_N)$ is the wave function renormalization constant, we obtain the following effective Lagrangian:

$$\mathcal{L}(x) = \partial_\mu S^\dagger(x) \partial^\mu S(x) - m_S^2 S^\dagger(x) S(x). \quad (6)$$

The mass of the field $S(x)$ is equal to

$$m_S = Z^{-1/2} \sqrt{4m_N^2 - \frac{g_S^2}{4\pi^2} J_1(m_N)} = 2m_N - \frac{g_S^2}{4\pi^2} m_N J_2(m_N), \quad (7)$$

where we have used the relation (5).

Following Ref.[3] we define the binding energy of the np pair in the spin-singlet state, described by the local interpolating field $S(x)$. It reads

$$\varepsilon_S = \frac{g_S^2}{4\pi^2} m_N J_2(m_N) = \frac{g_V^2}{18\pi^2} \frac{\Lambda_D^3}{m_N^2} = (0.079 \pm 0.012) \text{ MeV}, \quad (8)$$

where ± 0.012 is a theoretical accuracy of the approach [3]–[7].

The theoretical value of the binding energy $\varepsilon_S = (0.079 \pm 0.012)$ MeV agrees well with the binding energy $\varepsilon_{1S_0} = 0.074$ MeV, defined by the experimental S-wave scattering length of np scattering in the spin-singlet state.

⁴The coupling constant $g_V^2 = 2\pi^2 m_N^2 Q_D = (11.3)^2$, where $Q_D = 0.286 \text{ fm}^2$ is the electric quadrupole moment of the deuteron, is responsible for the existence of the deuteron as a Cooper np -pair [3].

⁵Such a relation can be easily obtained in the one-pion exchange approximation [5].

Using Eq.(8) we can compute the S-wave scattering length of np scattering in the spin-singlet state. Following Landau and Lifshitz [1], we obtain

$$a_S = -\frac{1}{\sqrt{\varepsilon_S m_N}} = (-23.00 \pm 1.74) \text{ fm.} \quad (9)$$

The theoretical value of a_S agrees well with the experimental one $a_{np} = (-23.75 \pm 0.01) \text{ fm}$.

Thus, we have shown that, as it has been suggested by Ericson [9], the np pair in the spin-singlet state can be treated as a Cooper np -pair with mass to be very close to the threshold mass $2m_N$. The difference ε_S between the threshold mass and the mass of the Cooper np -pair mass can be identified with the binding energy of the *virtual level* [1]. The theoretical value $\varepsilon_S = (0.079 \pm 0.012) \text{ MeV}$ agrees well with the the binding energy $\varepsilon_{1S_0} = 0.074 \text{ MeV}$ of the *virtual level* defined by the experimental value $a_{np} = (-23.75 \pm 0.01) \text{ fm}$ of the S-wave scattering length of np scattering in the spin-singlet state.

We are grateful to Torleif Ericson for the formulation of the problem and numerous helpful discussions. Natalia Troitskaya and Andrei Ivanov thank Torleif Ericson and Magda Ericson and TH-division of CERN for kind hospitality extended to them during the period of their stay at CERN, when this work was begun.

References

- [1] L. D. Landau and E. M. Lifshitz, in *QUANTUM MECHANICS, Non-relativistic Theory*, Second Edition, Pergamon Press, Oxford, 1965.
- [2] T. E. O. Ericson and W. Weise, in *PIONS AND NUCLEI*, Clarendon Press, Oxford, 1988; M. M. Nagels *et al.*, Nucl. Phys. B **147**, 189 (1979).
- [3] A. N. Ivanov, N. I. Troitskaya, H. Oberhummer, and M. Faber, Eur. Phys. J. A **7**, 519 (2000), nucl-th/0006049.
- [4] A. N. Ivanov, N. I. Troitskaya, H. Oberhummer, and M. Faber, Eur. Phys. J. A **8**, 129 (2000), nucl-th/0006050.
- [5] A. N. Ivanov, N. I. Troitskaya, H. Oberhummer, and M. Faber, Eur. Phys. J. A **8**, 223 (2000), nucl-th/0006051.
- [6] A. N. Ivanov, V. A. Ivanova, H. Oberhummer, N. I. Troitskaya and M. Faber, Eur. Phys. J. A **12**, 87 (2001), nucl-th/0108067.
- [7] A. N. Ivanov, M. Cargnelli, M. Faber, H. Fuhrmann, V. A. Ivanova, J. Marton, N. I. Troitskaya, and J. Zmeskal, nucl-th/0406081.
- [8] T. E. O. Ericson and M. Rosa-Clot, Phys. Lett. B **110**, 193 (1982); T. E. O. Ericson, Nuovo Cimento, **76**, 277 (1983); T. E. O. Ericson and M. Rosa-Clot, Nucl. Phys. A **405**, 497 (1983); M. Kamionkowski and J. N. Bahcall, Astrophys. J. **420**, 884 (1994).
- [9] T. E. O. Ericson, (private communication), CERN TH-Division, July of 2004.
- [10] T. E. O. Ericson and G. A. Miller, Phys. Lett. B **132**, 32 (1983).